

**PLASTIC SURGERY PRE OP ORDERS**

**Abdominoplasty, all Breast Reconstructions or Augmentation, and Mammoplasty Reductions**

- Cefazolin (KEFZOL) 1 gm IVPB 30 minutes Pre-Op (patient weight <50kg) – Administer within 1 hour of surgery start time.
- Cefazolin (KEFZOL) 2 gm IVPB (patient weight 50-120kg) – Administer within 1 hour of surgery start time.
- Cefazolin (KEFZOL) 3 gm IVPB (weights >120kg) – Administer within 1 hour of surgery start time.

**If life-threatening angioedema or anaphylaxis to cephalosporin or penicillin:**

- Vancomycin 1 gm IVPB (weight<100 kg) – Administer within 90 minutes of surgery start time to infuse over at least 90 minutes.
- Vancomycin 1500 mg gm IVPB (weight ≥100kg) – Administer within 2 hours of surgery start time to infuse over at least 120 minutes

**VTE PROPHYLAXIS:**

- Heparin 5000 units Subcut x1 dose @ \_\_\_\_\_ am/pm
- Apply and Turn **ON** Sequential Pneumatic Compression (SCD) Device  
**(PRIOR TO INDUCTION OF ANESTHESIA)**
- IV Acetaminophen (OFirmev) 1000mg / 100ml Pre-Op within 1 hour of surgery

\_\_\_\_\_  
Physician Signature

\_\_\_\_\_  
Date (required)

\_\_\_\_\_  
Time (required)



PT  
MR.#/P  
DR.