

McLaren Print System Order

Order No: 43587
 Order Date: 2019-03-14
 User: Judy Fago
 Phone: 586-493-3610

Ship Location: Gratiot Medical Bldg, Multi Specialty, Attn: Judy Fago
 36500 Gratiot, Suite 102
 Clinton Twp, MI 48035

Brochures
 Quantity: 200
 Paragon Dept No: 60330
 Dept Name: Multi
 Company Number: 260

Order Total Price: 39.60

Item Number: MO-74
 Item Description: Opioid Factsheet for Patients
 Revision Date: 3/2019
 Print:
 Paper:
 Size:
 Fold:
 Finish:
 Drill:
 Misc Info: ds; color; no bleed; 32#

PRESCRIPTION OPIOIDS: WHAT YOU NEED TO KNOW

Prescription opioids can be used to help relieve moderate-to-severe pain and are often prescribed following a surgery or injury, or for certain health conditions. These medications can be an important part of treatment but also come with serious risks. It is important to work with your health care provider to make sure you are getting the safest, most effective care.

WHAT ARE THE RISKS AND SIDE EFFECTS OF OPIOID USE?

Prescription opioids carry serious risks of addiction and overdose, especially with prolonged use. An opioid overdose, often marked by slowed breathing, can cause sudden death. The use of prescription opioids can have a number of side effects as well, even when taken as directed.

As many as 1 in 4 PEOPLE* receiving prescription opioids long term in a primary care setting struggle with addiction.

- Tolerance—meaning you might need to take more of a medication for the same pain relief
- Physical dependence—meaning you have symptoms of withdrawal when a medication is stopped
- Increased sensitivity to pain
- Constipation
- Nausea, vomiting, and dry mouth
- Sleepiness and dizziness
- Confusion
- Depression
- Low levels of testosterone that can result in less sex drive, energy and strength
- Itching and sweating

RISKS ARE GREATER WITH:

- History of drug misuse, substance use disorder, or overdose
- Mental health conditions (such as depression)
- Deep sleep
- Older age (65 years or older)
- Pregnancy

Avoid alcohol while taking prescription opioids. Also, unless specifically advised by your health care provider, medications to avoid include:

- Benzodiazepines (such as Xanax or Valium)
- Muscle relaxants (such as Soma or Flexeril)
- Sedatives (such as Ambien or Lunesta)
- Other prescription opioids

© 2019 CDC

Spec Info: