

McLaren Print System Order

Order No: 46011
Order Date: 2019-06-05
User: Lisa Jaeger
Phone: 586-493-2236

Ship Location: McLaren Macomb Attn: Jacqueline Phillips Respiratory Care
1000 Harrington Blvd.
Mount Clemens, MI 48043

Brochures
Quantity: 500
Paragon Dept No: 40110
Dept Name: Respiratory Care & Neurology Serv
Company Number: 60

Order Total Price: 20000.00

Item Number: M-2662
Item Description: Respiratory Medications Educational Material
Revision Date: 1/2019
Print:
Paper:
Size:
Fold:
Finish:
Drill:
Misc Info: 100; ds; color; 11x17 folded in half; 70#

Spec Info:

Picture	Brand Name	Generic Name	Directions for Use
	Albuterol (Sandoz)	Albuterol	Use inhalations every 4 to 6 hours, as needed or scheduled. Inhalations every 4 to 6 hours, as needed or scheduled.
	Formoterol (Sandoz)	Formoterol	Use inhalations every 12 hours, as needed or scheduled. Inhalations every 12 hours, as needed or scheduled.
	Budesonide (Sandoz)	Budesonide	Use inhalations every 2 to 4 hours, as needed or scheduled. Inhalations every 2 to 4 hours, as needed or scheduled.
	Budesonide (Sandoz)	Budesonide	Use inhalations every 2 to 4 hours, as needed or scheduled. Inhalations every 2 to 4 hours, as needed or scheduled.
	Budesonide (Sandoz)	Budesonide	Use inhalations every 2 to 4 hours, as needed or scheduled. Inhalations every 2 to 4 hours, as needed or scheduled.
	Budesonide (Sandoz)	Budesonide	Use inhalations every 2 to 4 hours, as needed or scheduled. Inhalations every 2 to 4 hours, as needed or scheduled.
	Budesonide (Sandoz)	Budesonide	Use inhalations every 2 to 4 hours, as needed or scheduled. Inhalations every 2 to 4 hours, as needed or scheduled.
	Budesonide (Sandoz)	Budesonide	Use inhalations every 2 to 4 hours, as needed or scheduled. Inhalations every 2 to 4 hours, as needed or scheduled.
	Budesonide (Sandoz)	Budesonide	Use inhalations every 2 to 4 hours, as needed or scheduled. Inhalations every 2 to 4 hours, as needed or scheduled.
	Budesonide (Sandoz)	Budesonide	Use inhalations every 2 to 4 hours, as needed or scheduled. Inhalations every 2 to 4 hours, as needed or scheduled.

See Educational Purposes Only

My Scheduled Medication	My Scheduled Medication	My Scheduled Medication

- General instructions for most inhaled medications:**
- Read the specific instructions provided for your medication(s). Talk to your pharmacist or doctor if you are unsure of how to use any medications.
 - Check the date printed to let you know when your inhaler needs to be refilled. Call your pharmacy for a refill when you have a week's supply of medication left.
 - Breathe out away from the inhaler.
 - Place the mouthpiece in your mouth and close your lips around it to form a seal.
 - Inhale following instructions for your specific inhaler.
 - Hold your breath for 5-10 seconds or as long as you are able.
 - Breathe out away from the inhaler through your nose.
 - Rinse out your mouth following each use. This is especially important for the inhaled combination medications.

The information contained herein may be updated or modified without notice. All trademarks are the property of their respective owners. © 2019 McLaren Macomb. All rights reserved. For more information, visit www.mclaren.com.