

McLaren Print System Order

Order No: 47822 Reprint Previous Order No: 17644
Order Date: 2019-08-19
User: Kim Brass
Phone: 8103001855

Ship Location: MMG Marysville Internal Med
1750 Busha Hwy
Marysville, MI 48040

Forms

Quantity: 100
Paragon Dept No: 66300
Dept Name: 66300
Company Number: 810

Order Total Price: 0.00

Item Number: MM-316
Item Description: Chickenpox (Varicella)
Revision Date: 3/2008
Print: 2 sided black and white
Paper: 20# White Text
Size: 8.5 x 11
Fold:
Finish: None
Drill: None
Misc Info: ds; black and white

VACCINE INFORMATION STATEMENT

Chickenpox Vaccine

What You Need to Know

Read this Vaccine Information Statement (VIS) carefully before you get this vaccine. There may be information on this vaccine that is not on this VIS. Talk to your healthcare provider.

1 Why get vaccinated?

Chickenpox (also called varicella) is a common childhood disease. It is usually mild, but it can be serious, especially in young infants and adults.

- It causes a rash, itching, fever, and tiredness.
- It can lead to severe skin infection, scars, pneumonia, brain damage, or death.
- The chickenpox virus can be spread from person to person through the air, or by contact with fluid from chickenpox blisters.
- A person who has had chickenpox can get a painful rash called shingles years later.
- Before the vaccine, about 11,000 people were hospitalized for chickenpox each year in the United States.
- Before the vaccine, about 100 people died each year as a result of chickenpox in the United States.

Chickenpox vaccine can prevent chickenpox.

Most people who get chickenpox vaccine will not get chickenpox. But if someone who has been vaccinated does get chickenpox, it is usually very mild. They will have fewer blisters, are less likely to have a fever, and will recover faster.

2 Who should get chickenpox vaccine and when?

Routine

Children who have never had chickenpox should get 2 doses of chickenpox vaccine at these ages:

1st Dose:	12-15 months of age
2nd Dose:	4-6 years of age (may be given earlier, if at least 3 months after the 1st dose)

People 12 years of age and older (who have never had chickenpox or received chickenpox vaccine) should get two doses at least 28 days apart.

3 Some people should not get chickenpox vaccine or should wait.

- People should not get chickenpox vaccine if they have ever had a life-threatening allergic reaction to a previous dose of chickenpox vaccine or to gelatin or the antibiotic neomycin.
- People who are moderately or severely ill at the time the shot is scheduled should usually wait until they recover before getting chickenpox vaccine.
- Pregnant women should wait to get chickenpox vaccine until after they have given birth. Women should not get pregnant for 1 month after getting chickenpox vaccine.
- Some people should check with their doctor about whether they should get chickenpox vaccine, including anyone who:
 - Has HIV/AIDS or another disease that affects the immune system
 - Is being treated with drugs that affect the immune system, such as steroids, for 2 weeks or longer
 - Has any kind of cancer
 - Is getting cancer treatment with radiation or drugs
- People who recently had a transfusion or were given other blood products should ask their doctor when they may get chickenpox vaccine.

Ask your doctor for more information.

U.S. Department of Health and Human Services
Centers for Disease Control and Prevention