

McLaren Print System Order

Order No: 48052
Order Date: 2019-08-27
User: susan jackson
Phone: 8103425370

Ship Location: McLaren Pulmonary Rehab
G 3230 Beecher Road lower level
Flint, Michigan 48532,

Brochures
Quantity: 100
Paragon Dept No: 40110
Dept Name: Pulmonary Rehab
Company Number: 60

Order Total Price: 8.76

Item Number: M-2651
Item Description: Exercise Guidelines
Revision Date: 6/2016
Print: 2 sided black and white
Paper: 20# White Text
Size: 8.5 x 11
Fold:
Finish: None
Drill:
Misc Info: 4 pages



Exercise Guidelines

Signs and symptoms of overexertion

Stop exercising if you experience an event that needs to be reported to a staff person such as:

- Extreme shortness of breath
- Chest pain or discomfort
- Nausea
- Leg pain or muscle cramps
- Dizziness
- Unusual fatigue
- Heart rate above "target heart rate range"

Additional Diabetic Exercising Guidelines

- You cannot exercise if it has been less than 1 hour since a meal.
- You cannot exercise if your blood sugar is over 250 or if your blood sugar is less than 80.

STRETCHING OUT TO BREATHE EASIER

When the lungs have to struggle to work properly, people start to use the muscles in the chest, neck, and shoulders to assist lung function, explains Dr. Schachter. "This tends to tighten up the upper body and rob strength from the other functions of these large muscles," he says. "This can be felt when a person raises his or her hands to hang curtains or use a hair dryer. If these and similar activities cause a person to feel short of breath, it is because the shoulder muscles (and other muscles) are already working hard to help the lungs." Flexibility exercises can help stretch, or loosen up, the muscles that have become involved in assisted breathing.

Spec Info: