

McLaren Print System Order

Order No: 61342 Reprint Previous Order No: 17839
 Order Date: 2021-03-28
 User: Theda Simmonds
 Phone: 989-393-2857

Ship Location: McLaren Occupational and Convenient Care - Bay City Occ
 4 Columbus Ave Ste 140
 Bay City, MI 48708

Forms

Quantity: 1000
 Paragon Dept No: 56052
 Dept Name: Occupational Convenient Care
 Company Number: 810

Order Total Price: 0.00

Item Number: MM-333
 Item Description: Td (Tetanus, Diphtheria)
 Revision Date: 4/11/2017
 Print: 2 sided black and white
 Paper: 20# White Text
 Size: 8.5 x 11
 Fold:
 Finish: None
 Drill: None
 Misc Info: ds; black and white

VACCINE INFORMATION STATEMENT		
<h2>Td Vaccine</h2> <h3>What You Need to Know</h3> <div style="border: 1px solid black; padding: 5px;"> <p>1 Why get vaccinated?</p> <p>Tetanus and diphtheria are very serious diseases. They are rare in the United States today, but people who do become infected often have severe complications. Td vaccine is used to protect adolescents and adults from both of these diseases.</p> <p>Both tetanus and diphtheria are infections caused by bacteria. Diphtheria spreads from person to person through coughing or sneezing. Tetanus-causing bacteria enter the body through cuts, scratches, or wounds.</p> <p>TETANUS (lockjaw) causes painful muscle tightening and stiffness, usually all over the body.</p> <ul style="list-style-type: none"> It can lead to tightening of muscles in the head and neck so you can't open your mouth, swallow, or sometimes even breathe. Tetanus kills about 1 out of every 10 people who are infected even after receiving the best medical care. <p>DIPHTHERIA can cause a thick coating to form in the back of the throat.</p> <ul style="list-style-type: none"> It can lead to breathing problems, paralysis, heart failure, and death. <p>Before vaccines, as many as 200,000 cases of diphtheria and hundreds of cases of tetanus were reported in the United States each year. Since vaccination began, reports of cases for both diseases have dropped by about 99%.</p> </div> <div style="border: 1px solid black; padding: 5px;"> <p>2 Td vaccine</p> <p>Td vaccine can protect adolescents and adults from tetanus and diphtheria. Td is usually given as a booster shot every 10 years but it can also be given earlier after a scrape and dirty wound or bite.</p> <p>Another vaccine, called Tdap, which protects against pertussis in addition to tetanus and diphtheria, is sometimes recommended instead of Td vaccine.</p> <p>Your doctor or the person giving you the vaccine can give you more information.</p> <p>Td may safely be given at the same time as other vaccines.</p> </div>	<p>(Tetanus and Diphtheria)</p> <p><small>More vaccine information statements are available in Spanish and other languages. See www.cdc.gov/vaccines. Steps to information from vaccine vials: diphtheria or tetanus + or another other illness. Visit www.cdc.gov/vaccines.</small></p>	<div style="border: 1px solid black; padding: 5px;"> <p>3 Some people should not get this vaccine</p> <ul style="list-style-type: none"> A person who has ever had a life-threatening allergic reaction after a previous dose of any tetanus or diphtheria containing vaccine, OR has a severe allergy to any part of this vaccine, should not get Td vaccine. Tell the person giving the vaccine about any severe allergies. Talk to your doctor if you: <ul style="list-style-type: none"> had severe pain or swelling after any vaccine containing diphtheria or tetanus, ever had a condition called Guillain Barre Syndrome (GBS), aren't feeling well on the day the shot is scheduled. </div> <div style="border: 1px solid black; padding: 5px;"> <p>4 Risks of a vaccine reaction</p> <p>With any medicine, including vaccines, there is a chance of side effects. These are usually mild and go away on their own. Serious reactions are also possible but are rare.</p> <p>Most people who get Td vaccine do not have any problems with it.</p> <p>Mild Problems following Td vaccine: (Did not interfere with activities):</p> <ul style="list-style-type: none"> Pain where the shot was given (about 8 people in 10) Redness or swelling where the shot was given (about 1 person in 4) Mild fever (rare) Headache (about 1 person in 4) Tiredness (about 1 person in 4) <p>Moderate Problems following Td vaccine: (Interfered with activities, but did not require medical attention):</p> <ul style="list-style-type: none"> Fever over 102°F (rare) <p>Serious Problems following Td vaccine: (Could not perform usual activities, required medical attention):</p> <ul style="list-style-type: none"> Swelling, severe pain, bleeding and/or redness in the arm where the shot was given (rare). </div>

