

McLaren Print System Order

Order No: 61343 Reprint Previous Order No: 17832
 Order Date: 2021-03-28
 User: Theda Simmonds
 Phone: 989-393-2857

Ship Location: McLaren Occupational and Convenient Care - Bay City Occ
 4 Columbus Ave Ste 140
 Bay City, MI 48708

Forms

Quantity: 1000
 Paragon Dept No: 56052
 Dept Name: Occupational Convenient Care
 Company Number: 810

Order Total Price: 0.00

Item Number: MM-326
 Item Description: MMR (Measles, Mumps, Rubella)
 Revision Date: 10/2020
 Print: 2 sided black and white
 Paper: 20# White Text
 Size: 8.5 x 11
 Fold: None
 Finish: None
 Drill: None
 Misc Info: ds; black and white

VACCINE INFORMATION STATEMENT

MMR Vaccine (Measles, Mumps, and Rubella): What You Need to Know

Many vaccine information statements are available in Spanish and other languages. For more information, visit www.cdc.gov/vaccines/imz/parents.
 Report information about vaccine safety concerns to the Vaccine Adverse Event Reporting System (VAERS) at www.vaers.hhs.gov.

1 Why get vaccinated?

MMR vaccine can prevent measles, mumps, and rubella.

- **MEASLES (M)** can cause fever, cough, runny nose, and red, watery eyes, commonly followed by a rash that covers the whole body. It can lead to seizures (often associated with fever), ear infections, diarrhea, and pneumonia. Rarely, measles can cause brain damage or death.
- **MUMPS (M)** can cause fever, headache, muscle aches, tiredness, loss of appetite, and swollen and tender salivary glands under the ears. It can lead to deafness, swelling of the brain and/or spinal cord covering, painful swelling of the testicles or ovaries, and, very rarely, death.
- **RUBELLA (R)** can cause fever, sore throat, rash, headache, and eye irritation. It can cause arthritis in up to half of teenage and adult women. If a woman gets rubella while she is pregnant, she could have a miscarriage or her baby could be born with serious birth defects.

Most people who are vaccinated with MMR will be protected for life. Vaccines and high rates of vaccination have made these diseases much less common in the United States.

2 MMR vaccine

Children need 2 doses of MMR vaccine, usually:

- First dose at 12 through 15 months of age
- Second dose at 4 through 6 years of age

Infants who will be traveling outside the United States when they are between 6 and 11 months of age should get a dose of MMR vaccine before travel. The child should still get 2 doses at the recommended ages for long-lasting protection.

Older children, adolescents, and adults also need 1 or 2 doses of MMR vaccine if they are not already immune to measles, mumps, and rubella. Your

3 Talk with your health care provider

Tell your vaccine provider if the person getting the vaccine:

- Has had an allergic reaction after a previous dose of MMR or MMRV vaccine, or has any severe, life-threatening allergies.
- Is pregnant, or thinks she might be pregnant.
- Has a weakened immune system, or has a parent, brother, or sister with a history of hereditary or congenital immune system problems.
- Has ever had a condition that makes him or her bruise or bleed easily.
- Has recently had a blood transfusion or received other blood products.
- Has tuberculosis.
- Has gotten any other vaccines in the past 4 weeks.

In some cases, your health care provider may decide to postpone MMR vaccination to a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting MMR vaccine.

Your health-care provider can give you more information.

U.S. Department of Health and Human Services
Centers for Disease Control and Prevention